

Supplemental Table 1. Assignment of individuals into parental and hybrid classes. Analyses include the linked data in STRUCTURE (K = 5; L16), unlinked data (haplotypes) in STRUCTURE (K = 2; this paper), analysis of gene trees (Tree), and unlinked data in NEWHYBRIDS (NH). For the last analysis, some individuals were not assigned to any class with confidence (Unresolved). Localities correspond with Table 1, Figure 1, and Figure 3.

Locality	n	<i>P. cinereus</i>				<i>P. electromorphus</i>				Hybrids			Unresolved	
		Linked	Unlinked	Tree	NH	Linked	Unlinked	Tree2	NH	Linked	Unlinked	Tree2	NH	NH
1	2	0	2	2	2	0	0	0	0	2	0	0	0	0
2	5	0	5	5	4	0	0	0	0	5	0	0	0	1
3	2	0	2	1	2	0	0	0	0	2	0	1	0	0
4	4	1	4	3	2	0	0	0	0	3	0	1	0	2
5	3	1	2	3	2	0	0	0	0	2	1	0	0	1
6	4	0	0	0	0	1	3	3	2	3	1	1	0	2
7	36	10	18	14	10	8	12	11	9	18	6	11	3	14
8	3	0	1	1	1	1	2	1	2	2	0	1	0	0
9	5	3	5	4	2	0	0	0	0	2	0	1	0	3
10	6	3	3	3	3	1	3	3	3	2	0	0	0	0
11	6	0	5	4	4	0	1	0	0	7	0	2	0	2
12	8	0	0	0	0	8	8	8	7	0	0	0	0	1
13	5	0	0	0	0	5	5	5	3	0	0	0	0	2